

CpDNA-based species identification and phylogeography: application to a complex of African tropical tree species

Duminil J.¹, Heuertz M.^{1,2}, Doucet J.-L.³, Bourland N.³, Cruaud C.⁴;
Doumenge C.⁵, Navascués M.^{6,7} & Hardy O.¹.

¹ Service Evolution Biologique et Ecologie, ULB, 50 Av. F. Roosevelt, B-1050 Bruxelles, Belgique

² Department of Biodiversity and Conservation, Real Jardín Botánico, CSIC, Plaza de Murillo, 2, E-28014 Madrid, Spain

³ Laboratoire de Foresterie des Régions tropicales et subtropicales, Unité de Gestion des Ressources forestières et des Milieux naturels, Gembloux Agro-Bio Tech, Université de Liège, Passage des Déportés 2, B-5030 Gembloux, Belgique

⁴ CEA, DSV, Institut de Génomique, Genoscope, 2 rue Gaston Crémieux, CP5706, F-91057 Evry, France

⁵ Unité Biens et Services des Ecosystèmes Forestiers, CIRAD, TA C-105/D, Campus international de Baillarguet, F-34398 Montpellier Cedex 5, France

⁶ Équipe Éco-évolution Mathématique. CNRS UMR 7625 Écologie et Évolution. Université Pierre et Marie Curie, École Normale Supérieure, UMR 7625, 46 rue d'Ulm, F-75230 Paris Cedex 05, France

⁷ INRA, UMR CBGP (INRA/IRD/Cirad/Montpellier SupAgro), Campus International de Baillarguet, CS 30016, F-34988 Montferrier-sur-Lez Cedex, France

e-mail pour correspondance : jduminil@ulb.ac.be

Mots-clefs : pleistocene forest refuges, genetic structure, human influence, species delimitation, African rainforest

Résumé : Despite their importance as a hotspot of biodiversity, the history of the African tropical lowland rainforests is poorly known. In particular the respective influence of past climatic factors, environmental heterogeneity and physical barriers on the demographic history of plant species still have to be studied. Molecular approaches that characterise the distribution of genetic diversity in plant species and try to interpret it in the light of past historical events are promising in this respect. We examine here the distribution of genetic diversity at maternally-inherited markers (chloroplast DNA) of two tropical timber tree species, *Erythrophleum ivorense* and *Erythrophleum suaveolens* (Leguminosae-Caesalpinioideae), in their Lower Guinea distribution (essentially Gabon and Cameroon). As for many tropical sister-taxa, species identification is almost impossible in the field when flowers or fruits are not available. Using a blind-sampling approach and confronting molecular and morphological markers it was possible to attribute sampled individuals to one or the other species. The two species appear to have different ecological requirements and are rarely found in sympatry. A phylogeographic signal was found for *E. suaveolens* and, for both species, higher levels of diversity were observed in the South than in the North of the Lower Guinea domain. A recent period of forest perturbation, potentially human-induced, has probably favoured the expansion of these two light-demanding species. An observed genetic signal of demographic decline and the poor recruitment of the species can be interpreted as consequences of the natural succession of the forest formations.